

## **PHEASANT AND PARTRIDGE WEIGHT AND WATER CHART**

PLEASE NOTE: THESE VALUES ARE ALL APPROXIMATE, AND WILL FLUCTUATE THROUGHOUT THE SEASON, DEPENDANT MAINLY ON WEATHER.

AGE (WEEKS)	PHEASANT 1000 birds			RED-LEGGED PARTRIDGE 1000 birds			GREY PARTRIDGE 1000 birds		
	Water Consumption		Weight	Water Consumption		Weight	Water Consumption		Weight
	gallons	litres	Kg	gallons	litres	Kg	gallons	litres	Kg
0(day old)	2	9	20	1	4.5	12	0.7	3.2	7
1	3	13	50	1.5	6.8	17	1	4.5	11
2	4.5	20	80	2.3	10.3	30	1.5	6.8	18
3	6	27	120	3	13.5	45	2	9	28
4	9	40	200	4.5	20	70	3	13.5	45
5	12	54	300	6	27	100	4	18	62
6	13	60	380	6.5	30	140	4.3	19.4	86
7	14	64	450	7	32	190	4.7	21	115
8	15	68	550	7.5	34	230	5	22.5	140
9	16	73	650	8	36	280	5.3	24	170
10	16	73	750	8	36	330	5.5	25	210
12	17	78	900	8.5	38	400	5.7	26	250
ADULT									
Male	27	120	1400	11.5	52	540	7	32	390
Female	21	95	1100	10.2	46	480	6.5	30	330

## Baycox / Toltra-K / Zorabel for Gamebirds.

Dose (ml) = 0.28 X liveweight/Kg of birds.

Age (weeks)	PHEASANT	PARTRIDGE
	ml/ litres Drinking water	ml/ litres Drinking water
1	n/a	n/a
2	1.1	0.84
3	1.2	0.9
4	1.4	1
5	1.5	1
6	1.7	1.3
7	2	1.7
8	2.2	1.9
9	2.2	2.2
10	2.2	2.2
12	2.2	2.2
Adult	2.2	2.2

### Dosage Notes

Add 1 to 2.2 ml per litre according to the dosage chart opposite

**Note: over 2.2ml per litre the birds do not like the product so water consumption may decline. Better to run at 2.2ml per litre for 3-4 days**

No other water to be available during treatment

Make fresh twice daily

1 Gallon = 4.55 litres

40 Gallons = 182 litres

### Baycox Verdunner/ NaOH Water Additive (Buffer)

This helps keep the Baycox in solution in hard water and we may recommend it when prescribing treatment.

Add an hour before adding the Baycox, especially if water is from a borehole.

100ml treats 4000 litres of water.

So 5ml treats a 200 litre tank.

**Note – this is highly caustic when concentrated so wear gloves when handling.**

NB all consumption levels will be minimum to provide fresh each day, and one should allow for wastage.

The solution should be made up fresh twice daily in hardwater areas and warm weather.

**Treatment Time: 3 days**

**Withdrawal time (meat) 28 days**

NOTE: All charts are provided for the exclusive use of South Downs Veterinary Consultancy and no responsibility can be taken for use of these doses by others