

Soludox/ Hydrodoxx/Doxytib/Primox: Dosing guide.

(water)50mg/Kg

Product is 50% active ingredient by weight so multiply need by 2 to get gram equivalent by weight of product (which is the amount shown to administer in the chart)

Age (weeks)	PHEASANT		PARTRIDGE	
	Gram of product/1000 birds	g/100 litres Drinking water	g/1000 birds	g/100 litres Drinking water
1	5	40	1.7	25
2	8	40	3	30
3	12	45	4.5	35
4	20	50	7	35
5	30	55	10	37.5
6	38	65	14	46.5
7	45	70	19	60
8	55	80	23	70
9	65	90	28	80
10	75	105	33	92.5
12	90	115	40	105
Adult	140	115	55	110

Notes:

For ease of use, at 7 weeks old the Dose is 1 level 5ml teaspoon in 5 litres (1 gallon) of water

1 x 5ml teaspoon holds 3.6g of product.

In Hardwater areas twice the normal dosage can be required to keep the product in solution. So it is worth considering using water Acidity regulators such as Salkil for easier use

NB all consumption levels will be minimum to provide fresh each day, and one should allow for wastage. The solution should be made up fresh twice daily in hardwater areas and warm weather.

Treatment Time 5-7days

Withdrawal time (meat) 28 days

NOTE: All charts are provided for the exclusive use of South Downs Veterinary Consultancy and no responsibility can be taken for use of these doses by others

Karidox/Powdox Liquid: Dosing guide for Gamebirds.

(water)50mg/Kg

Product is 10% active ingredient by weight

Age (weeks)	PHEASANT		PARTRIDGE - RedL		PARTRIDGE - Grey	
	g/1000 birds Doxycyc	ml/100 litres Drinking water	g/1000 birds Doxycyc	ml/100 litres Drinking water	g/1000 birds Doxycyc	ml/100 litres Drinking water
1	2.5	192	0.85	125	0.55	122
2	4	192	1.5	146	0.9	132
3	6	222	2.25	167	1.4	155
4	10	250	3.5	175	2.25	167
5	15	277	5	185	3.1	172
6	19	316	7	233	4.3	221
7	22.5	351	9.5	297	5.75	273
8	27.5	404	11.5	338	7	311
9	32.5	445	14	388	8.5	354
10	37.5	510	16.5	458	10.5	420
12	45	577	20	526	12.5	480
Adult	70	700	22.5	561	17.5	564

NB all consumption levels will be minimum to provide fresh each day, and one should allow for wastage. The solution should be made up fresh twice daily in hardwater areas and warm weather.

Treatment Time 5-7days

Withdrawal time (meat) 28 days

NOTE: All charts are provided for the exclusive use of South Downs Veterinary Consultancy and no responsibility can be taken for use of these doses by others